

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

8 e 7.

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C07C 381/00		A2	(11) International Publication Number: WO 92/17445
			(43) International Publication Date: 15 October 1992 (15.10.92)
(21) International Application Number: PCT/US92/02560		(74) Agents: GOLDSTEIN, Jorge, A. et al.; Sterne, Kessler, Goldstein & Fox, 1225 Connecticut Avenue, N.W., Suite 300, Washington, DC 20036 (US).	
(22) International Filing Date: 30 March 1992 (30.03.92)			
(30) Priority data: 676,691 29 March 1991 (29.03.91) US 804,665 11 December 1991 (11.12.91) US		(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), MC (European patent), NL (European patent), SE (European patent).	
(71) Applicant: BRIGHAM AND WOMEN'S HOSPITAL [US/US]; 75 Francis Street, Boston, MA 02115 (US).		Published Without international search report and to be republished upon receipt of that report.	
(72) Inventors: STAMLER, Jonathan, S.; 220 Marlborough Street, #1, Boston, MA 02116 (US). LOSCALZO, Joseph; 50 Pacella Drive, Dedham, MA 02026 (US). SLIVKA, Adam; 33 Stoughton Street, Randolph, MA 02368 (US). SIMON, Daniel; 211 Dorset Road, Waban, MA 02168 (US). BROWN, Robert; 12 Farmhill Road, Natick, MA 01760 (US). DRAZEN, Jeffrey; 99 Lawson Road, Winchester, MA 01890 (US).			

(54) Title: S-NITROSTHIOLS AS SMOOTH MUSCLE RELAXANTS AND THERAPEUTIC USES THEREOF



(57) Abstract

S-nitrosothiols exert a potent relaxant effect, mediated by guanylate cyclase, upon non-vascular smooth muscle. Such types of smooth muscle include airway, gastrointestinal, bladder, uterine and corpus cavernosal. Thus, S-nitrosothiols may be used for the treatment or prevention of disorders associated with relaxation of smooth muscle, such as airway obstruction, and other respiratory disorders, bladder dysfunction, premature labor and impotence. Additionally, S-nitrosothiols may be used to alleviate smooth muscle contraction and spasm, and thus facilitate procedures involving diagnostic instrumentation, such as endoscopy, bronchoscopy, laparoscopy and cystoscopy.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
DE	Germany	MC	Monaco	TC	Togo
DK	Denmark			US	United States of America

- 2 -

nitroso-captopril, to relax non-vascular smooth muscle. Types of smooth muscle include airway, gastrointestinal, bladder uterine, and corpus cavernosum. The invention also relates to the use of S-nitrosothiols for the treatment or prevention of disorders which involve non-vascular smooth muscle, such as respiratory disorders, gastrointestinal disorders, urological dysfunction, impotence, uterine dysfunction or premature labor. The invention also relates to the use of S-nitrosothiols to ameliorate smooth muscle contraction or spasm and thus, facilitate diagnostic or therapeutic procedures, such as bronchoscopy, endoscopy, laparoscopy, and cystoscopy.

Brief Description of the Background Art

The endothelium secretes a vascular relaxing factor, known as endothelium-derived relaxing factor (EDRF), which has been identified as nitric oxide (NO), or a closely related derivative thereof. (Palmer *et al.*, *Nature* 327:524-526 (1987); Ignarro *et al.*, *Proc. Natl. Acad. Sci. USA* 84:9265-9269 (1987)). Under physiologic conditions, however, NO is exceedingly unstable, reacting essentially instantaneously with oxygen, superoxide anion, and redox metals (Lancaster *et al.*, *Proc. Natl. Acad. Sci. USA* 87:1223-1227 (1990); Ignarro *et al.*, *Circ. Res.* 65:1-21 (1989); and Gryglewski *et al.*, *Nature* 320:454-456 (1986)). This fact has lead to the supposition that, in order to exert its effect on vascular smooth muscle, NO must be stabilized *in vivo* in a form that preserves its biological activity.

S-nitrosothiols (RS-NO) are adducts that form readily under physiologic conditions from the reaction of NO with reduced low molecular weight thiols (Oae *et al.*, *Org. Prep. Proc. Int.* 15(3):165-198 (1983)). These compounds have half-lives that are significantly greater than that of NO and, like EDRF, possess vasorelaxant activity that is

- 3 -

mediated through activation of guanylate cyclase (Kowaluk *et al.*, *J. Pharmacol. Exp. Ther.* 256:1256-1264 (1990); Loscalzo *et al.*, *J. Pharmacol. Exp. Ther.* 249(3):726-729 (1989); and Ignarro *et al.*, *J. Pharmacol. Exp. Ther.* 218(3):739-749 (1981)).

5 The relaxant effect of S-nitrosothiols on blood vessels, and the mechanism by which this effect is exerted, is reasonably well understood in the art. However, the role of NO, or involvement of the guanylate cyclase pathway in non-vascular smooth muscle is not as clearly understood.

10 In the lung, pulmonary endothelial cells, macrophages and polymorphonuclear leukocytes are potential sources of NO or RS-NO. However, the role of NO and its metabolites in regulation of airway tone is not known, and the few available reports on the efficacy of NO and EDRF in relaxation of airway smooth muscle are conflicting (Shikano *et al.*, *J. Pharmacol. Exp. Ther.* 243:55-59 (1987); Shikano *et al.*, *Br. Journal Pharmacol.* 92:483-485 (1987)). Furthermore, in the lung, the high ambient concentrations of oxygen and other oxygen-free radicals predispose to rapid inactivation of NO (Furchgott R.F. *et al.*, *J. Endothelium-Derived Relaxing Factors and Nitric Oxide*; eds. Rubanyi G.M.,
15 pp. 8-21 (1990); Gryglewski, R.J. *et al.*, *Nature* 320:454-456 (1986)).

20 Non-vascular smooth muscle is present in numerous organ systems throughout the body, and has a vital role in the physiological function of these systems. For example, airway smooth muscle plays a critical role in constriction and dilation of bronchi. In the gastrointestinal tract, the
25 sphincter of Oddi, a smooth muscle connection between the bile duct and duodenum, provides tonic contraction which serves to prevent reflux of duodenal contents into the pancreatic and bile ducts, and promotes filling of the gall bladder. In addition, esophageal (sphincters and body), intestinal and colonic motility is regulated by smooth muscle. Smooth
30 muscle of the bladder body, bladder base, and proximal urethra plays an

- 4 -


important role in urological function, and erectile function is mediated by relaxation of corpus cavernosal smooth muscle.

In summary, the relaxation kinetics of non-vascular smooth muscle are very important in numerous physiological systems. Moreover, a variety of significant clinical disorders occur, which involve contraction, spasm, or failure to achieve the necessary relaxation of smooth muscle. Examples of such disorders include airway obstruction (i.e., asthma, bronchitis and emphysema), bladder dysfunction, gastrointestinal muscle spasm (i.e., irritable bowel syndrome, achylasia, dumping disorders), and impotence. Thus, a clinical need exists for pharmacological agents which can treat or prevent such disorders by inducing relaxation of the affected smooth muscle.

SUMMARY OF THE INVENTION

This invention is based on the discovery by the inventors that S-nitrosothiols exert a potent relaxant effect on non-vascular smooth muscle. This concept lead the inventors to the discovery that S-nitrosothiol compounds may be used as a therapeutic modality in disorders which involve smooth muscle relaxation.


The invention is directed to an S-nitrosothiol compound which has the formula:



wherein:

X equals 2 to 20.

The invention is also directed to an S-nitrosothiol compound which has the formula:




wherein:

- 5 -

X equals 2 to 20.

The invention is also directed to an S-nitrosothiol compound which has the formula:




5 wherein:

X equals 2 to 20 and Y is selected from the group consisting of fluoro, C₁-C₆ alkoxy, cyano, carboxamido, C₃-C₆ cycloalkyl, aralkoxy, C₂-C₆ alkylsulfinyl, arylthio, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbomoyl, C₁-C₆ N-alkylcarbomoyl, C₂-C₁₅ N,N-dialkylcarbomoyl, amino, 10 hydroxyl, carboxyl, hydrogen, nitro and aryl; wherein aryl includes benzyl, naphthyl, and anthracenyl groups.

The invention is also directed to the use of S-nitrosothiols for the treatment or prevention of disorders associated with relaxation of smooth muscle, such as airway obstruction, gastrointestinal spasm, bladder 15 dysfunction and impotence. The invention is also directed to the use of S-nitrosothiols to alleviate smooth muscle contraction and spasm, and thus facilitate procedures involving diagnostic instrumentation such as endoscopy and bronchoscopy.

In particular, this invention is directed to a method for relaxing 20 airway smooth muscle by administering a therapeutically effective amount of an S-nitrosothiol compound to an animal. The S-nitrosothiol compound may be selected from the group consisting of S-nitroso-N-acetylcysteine, S-nitroso-glutathione, S-nitroso-cysteine, S-nitroso-homocysteine, S-nitroso-penicillamine and S-nitroso-captopril. The S-nitrosothiol compound may be selected from the group consisting of a 25 compound having the formula:




wherein:

X equals 2 to 20.

- 6 -


The invention is also directed to an S-nitrosothiol compound which has the formula:



wherein:

5 X equals 2 to 20.

The invention is also directed to an S-nitrosothiol compound which has the formula:



wherein:

10 X equals 2 to 20 and Y is selected from the group consisting of fluoro, C₁-C₆ alkoxy, cyano, carboxamido, C₃-C₆ cycloalkyl, aralkoxy, C₂-C₆ alkylsulfinyl, arylthio, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbomoyl, C₁C₆ N-alkylcarbomoyl, C₂-C₁₅ N,N-dialkylcarbomoyl, amino, hydroxyl, carboxyl, hydrogen, nitro and aryl; wherein aryl includes benzyl, 15 naphthyl, and anthracenyl groups.

The invention is also directed to a method for treatment or prevention of respiratory disorders by administering a therapeutically effective amount of S-nitrosothiol compound to an animal. Respiratory disorders include obstructive lung disease, emphysema, asthma, bronchitis, 20 fibrosis, excessive mucus secretion, obstruction of air flow, and lung disorders resulting from post-surgical complications.

The invention is also directed to a method for relaxing gastrointestinal smooth muscle by administering a therapeutically effective amount of an S-nitrosothiol compound to an animal.

25 The invention is also directed to a method for ameliorating contraction or spasm of gastrointestinal smooth muscle associated with endoscopic procedures, by administering a therapeutically effective amount of an S-nitrosothiol compound to an animal.

- 7 -

The invention is also directed to a method for relaxing corpus cavernosum smooth muscle by administering a therapeutically effective amount of an S-nitrosothiol compound to an animal.

5 The invention is directed to a method for the treatment or prevention of impotence by administering a therapeutically effective amount of an S-nitrosothiol compound to an animal.

The invention is also directed to the administration of said S-nitrosothiol compounds for the methods of the invention, as part of the pharmaceutical composition comprising a pharmaceutically acceptable
10 carrier.

The invention is also directed to the methods of the invention wherein the pharmaceutical composition containing the S-nitrosothiol compound is administered to an animal by a route comprising oral, sublingual, intravenous, topical, intramuscular or intranasal delivery.

15 The invention is also directed to a method for relaxing bladder smooth muscle by administering a therapeutically effective amount of an S-nitrosothiol compound to an animal.

The invention is also directed to a method for relaxing uterine smooth muscle by administering a therapeutically effective amount of an
20 S-nitrosothiol compound to an animal.

Brief Description of the Figures

FIGURE 1: Inhibition of the Sphincter of Oddi by administration of S-nitroso-N-acetylcysteine.

FIGURE 2: Inhibition of duodenal motility by administration of
25 S-nitroso-N-acetylcysteine.

- 8 -

FIGURE 3: Side-by-side comparison of the relaxant effect of specific S-nitrosothiols on guinea pig tracheal muscle.

5 FIGURE 4: Dose-dependent relaxant effect of specific S-nitrosothiols on guinea pig tracheal muscle contracted with $3 \mu\text{M}$, as compared to the reactant and NO.

- 10 a: S-nitroso-glutathione
b: S-nitroso-cysteine
c: S-nitroso-homocysteine
d: S-nitroso-N-acetylcysteine
e: S-nitroso-penicillamine
f: S-nitroso-captopril

15 FIGURE 5: Relaxant activities of S-nitroso-N-acetylcysteine (A) and S-nitroso-captopril (B) determined against contractions induced by leukotriene D_4 , histamine and methacholine.

FIGURE 6: The course of relaxation induced by S-nitroso-N-acetylcysteine ($5 \times 10^{-6}\text{M}$) over 60 minutes.

20 FIGURE 7: The relaxation response to S-nitroso-glutathione in guinea pig airway (A) and rabbit aorta (B).

FIGURE 8: Tracheal relaxant effects of S-nitroso-N-acetylcysteine, isoproterenol, and theophylline.

- 9 -

FIGURE 9: Inhibition of tracheal relaxation to S-nitroso-N-acetylcysteine by hemoglobin and methylene blue.

FIGURE 10: Cyclic GMP determinations in tracheal rings incubated with 100 μ M S-nitroso-N-acetylcysteine.

5 FIGURE 11: Comparison between the relaxant effect of S-nitroso-glutathione and nitrite upon human tracheal smooth muscle.

10 FIGURE 12: Comparison between the relaxant effect of S-nitroso-glutathione and glutathione upon human tracheal smooth muscle.

FIGURE 13: Comparison between the relaxant effect of S-nitroso-N-acetylcysteine and N-acetylcysteine upon human tracheal smooth muscle.

15 FIGURE 14: Tracheal relaxant effects of theophylline, isoproterenol, S-nitroso-N-acetylcysteine, and S-nitroso-glutathione.

FIGURE 15: Cyclic GMP response to S-nitroso-N-acetylcysteine in human airways.

Description Of The Preferred EmbodimentsBackground

The invention is based on the discovery by the inventors that S-nitrosothiols relax non-vascular smooth muscle, and possess unique and different relaxant activities, kinetic properties and membrane permeability, and thus, may be used to treat or prevent disorders which involve non-vascular smooth muscle.

In one embodiment, the term "S-nitrosothiol" refers to a compound which is selected from the group consisting of S-nitroso-N-acetylcysteine, S-nitroso-glutathione, S-nitroso-cysteine, S-nitroso-homocysteine, S-nitroso-pantathoine derivatives, S-nitroso-penicillamine and S-nitroso-captopril.

In another embodiment the term "S-nitrosothiol" refers to particular novel S-nitrosothiol compounds synthesized by the inventors, for use as smooth muscle relaxants. The compounds represented by the general formula of $\text{CH}_3(\text{CH}_2)_x\text{SNO}$ are long carbon-chain lipophilic nitrosothiols. The compounds represented by the general formula of $\text{HS}(\text{CH}_2)_x\text{SNO}$ are S-nitrosodithiols, possessing an additional thiol group. The compounds represented by the general formula of $\text{ONS}(\text{CH}_2)_x\text{Y}$ are S-nitrosothiols which possess other functional groups, in addition to the thiol.

The invention is related to the discovery that S-nitrosothiol compounds relax non-vascular smooth muscle. As a result, these compounds may be used to treat or prevent those pathophysiologic conditions which result from, or involve, constriction of smooth muscle, or those which necessitate therapeutic intervention to achieve smooth muscle relaxation.

- 11 -

One embodiment of the invention relates to the administration of a therapeutically effective amount of an S-nitrosothiol to an animal to relax airway smooth muscle. The term "airway smooth muscle" refers to the smooth muscle lining the bronchi or tracheal region. The inventors
5 have demonstrated that S-nitrosothiols exert a potent relaxant effect upon airway smooth muscle.

As a result of this potent relaxant effect exerted by S-nitrosothiols, these compounds may be administered as therapeutic agents for the treatment or prevention of respiratory disorders.

10 The term "respiratory disorder" refers to any impairment of lung function which involves constriction of airways and changes in blood gas levels or lung function.

For example, airway obstruction constitutes a respiratory disorder which occurs as a result of acute pulmonary impairment or obstructive
15 lung disease. Severe airway obstruction may ultimately result in life-threatening respiratory failure. Airway obstruction occurs in patients with chronic obstructive lung diseases, such as emphysema and bronchitis. These patients often experience recurrent episodes of respiratory failure as a result of severe airway obstruction. Emphysema can result in
20 significant disability due to dyspnea, extreme restriction of physical activity, and mortality.

Airway obstruction also results from asthma, a disorder characterized by increased responsiveness of the tracheobronchial tree to various stimuli, and which leads to generalized airway constriction
25 manifested by dyspnea, cough and wheezing. Asthma sufferers often experience acute exacerbations of bronchoconstriction, which may be life-threatening.

Another obstructive lung disease, cystic fibrosis, results from abnormal exocrine gland function. Clinical manifestations include
30 excessive mucous secretion, hypertrophy of bronchial glands, infection.

- 12 -

and inflammatory and structural changes in the airways which lead to obstruction and ventilation-perfusion imbalance.

Acute respiratory failure may result not only from obstructive disease, but also as a consequence of airway constriction secondary to pneumonia, thromboembolism, left ventricular failure and pneumothorax. Acute respiratory failure may also result from ventilation-perfusion imbalance.

A critical component in the treatment of airway obstruction involves the use of pharmacologic agents to remove secretions and reverse airway constriction. The most commonly used bronchodilatory agents are beta-agonists, such as isoproterenol, given by inhalation or subcutaneous injection, and methylxanthines, such as theophylline, given orally or by infusion.

The margin of safety for theophylline administration is relatively narrow. The minimum therapeutic concentration in plasma is 6 to 10 $\mu\text{g/ml}$, and unacceptable symptoms of toxicity usually appear at or above 20 $\mu\text{g/ml}$. Still higher concentrations can lead to serious central nervous system toxicity, with long-term ingestion of theophylline being a predisposing factor in such toxicity. Moreover, because the clearance of theophylline is influenced by genetic, developmental and environmental factors to a significant degree, it is necessary to titrate the dosage cautiously against clinical observations of beneficial or toxic effects, with periodic determination of the concentration of the drug in plasma (Gilman A.G., *The Pharmacological Basis of Therapeutics*, Pergamon Press, New York, (1990)).

Isoproterenol, a non-selective β -agonist, produces cardiovascular side effects such as palpitations, sinus tachycardia and more serious arrhythmias. In addition, tolerance to this drug may result from overuse (Gilman A.G., *The Pharmacological Basis of Therapeutics*, Pergamon Press, New York, (1990)). This characteristic reduces its usefulness in patients

- 13 -

with chronic obstructive disease who rely heavily on frequent use of bronchodilators. It has now been demonstrated that β agonists may have long term deleterious effects which result in aggravation of asthma, and ultimately change the natural history of the disease. Consequently, the
5 American Thoracic Society no longer recommends treatment with β agonists as first line therapy in mild asthma (Expert Panel Recommendation, *New England Journal of Medicine*, 325:425-426 (1991)).

The use of S-nitrosothiols for the treatment of airway obstruction provides significant advantages over current methods of treatment. The
10 use of S-nitrosothiols eliminates the untoward side effects associated with β -agonists and methylxanthines.

Furthermore, because all current treatment methods act by way of cAMP, S-nitrosothiols satisfy the need for bronchodilators which act by way of cGMP. This is important because current evidence provided by
15 the inventors demonstrates a role for cyclic GMP in regulation of airway tone in humans (See Example 1). In addition, cyclic GMP agonists act synergistically with cyclic AMP agonists to provide bronchodilation, not obtainable by individual agents.

S-nitrosothiols also potently inhibit platelets and neutrophils, which
20 have been implicated in the pathogenesis of asthma. S-nitrosothiols provide nitric oxide in its biologically relevant form, which is critical because the bioactivity (relaxant activity) of nitric oxide in airways depends upon the form in which it is delivered.

Finally, the inventors have demonstrated that S-nitrosothiols
25 mediate the activity of vasoactive intestinal peptide (VIP), an important airway relaxant. This reinforces the importance of S-nitrosothiols in regulation of airway tone. Deficiency in the effect of VIP is a causal factor in the pathogenesis of asthma. Administration of S-nitrosothiols replenishes the mediator itself rather than a less biologically active
30 derivative.

- 14 -

S-nitrosothiols are also suitable for direct instillation into the lungs by bronchoscopic means. This topical administration permits titration of dose, eliminates the untoward side effects of systemic therapy, and enables the use of combination therapy, involving a topical S-nitrosothiol in conjunction with a systemic agent, in problematic cases. This topical therapy would also facilitate endoscopy by suppressing the cough reflex and associated bronchospasm.

An important component in the treatment of airway obstruction is the removal of airway mucous. Thus, airway obstruction often necessitates the administration of a mucolytic agent in conjunction with the bronchodilator. N-acetylcysteine, more commonly known as "Mucomist", is one such agent. S-nitroso-N-acetylcysteine possesses both mucolytic and bronchodilator capabilities.

With respect to combined bronchodilator-mucolytic agents, the mucolytic activity of the compound depends upon the amount of thiol which is preserved after NO delivery. Thus, S-nitrosothiol compounds which contain more than one thiol (dithiol compounds) are particularly suitable for achieving mucolysis. In addition, the long-chain lipophilic S-nitrosothiols which contain more than one thiol are advantageous as mucolytic agents because they have a free thiol, and their lipophilic property facilitates penetration of the compound into the lipid portion responsible for the tenacious viscosity of mucous.

In addition to the treatment or prevention of respiratory disorders, S-nitrosothiols may also be used to facilitate diagnostic and therapeutic bronchoscopy. The term "bronchoscopy" refers to the procedure in which a flexible fiberoptic, or rigid bronchoscope is introduced into the tracheobronchial tree for the purpose of bronchial visualization, lung biopsy or brushings, aspiration of secretions, and delivery of pharmacological agents.

- 15 -

A complication of bronchoscopy, and thus an impediment to the successful completion of the procedure, is bronchospasm. Patients with a prior history of bronchospasm are particularly at risk for acute enhancement of spasm. Thus, S-nitrosothiols may also be used to relax
5 airway smooth muscle and eliminate bronchoscopy-induced bronchospasm.

Another embodiment of the invention relates to the administration of a therapeutically effective amount of an S-nitrosothiol compound to an animal to relax gastrointestinal smooth muscle. The term "gastrointestinal smooth muscle" refers to smooth muscle which is contained in all areas
10 of the gastrointestinal tract. Such areas include, but are not limited to, the esophagus, duodenum, sphincter of Oddi, biliary tract, ileum, sigmoid colon, pancreatic duct and common bile duct. S-nitrosothiols may be used for the treatment or prevention of gastrointestinal disorders. Disorders of the gastrointestinal tract include achylasia (spasm of the lower
15 esophageal sphincter), diarrhea, dumping syndrome, and irritable bowel.

An additional embodiment of the invention relates to the administration of S-nitrosothiols to alleviate contraction or spasm of gastrointestinal smooth muscle, and thus facilitate successful completion of endoscopic procedures. Contraction or spasm of gastrointestinal
20 smooth muscle imposes a technical obstacle which must frequently be overcome in order to enable the clinician to successfully perform endoscopic procedures.

The term "endoscopic procedures" refers to those diagnostic procedures which utilize an instrument which is introduced into the
25 gastrointestinal tract to provide direct visualization of the gastrointestinal tract, for examination and therapeutic purposes. Such purposes include direct visualization, biopsy, access to the common bile duct, fluid aspiration and removal of foreign bodies, polyps, and other lesions. An example of a particular endoscopic procedure is esophagogastroduodenoscopy, which is utilized for examination of the esophageal lumen.
30

- 16 -

stomach and duodenum. Another example, endoscopic retrograde cholangiopancreatography (ERCP), enables visualization of the pancreatic duct, common bile duct and the entire biliary tract, including the gall bladder. Further examples of endoscopic procedures are colonoscopy and sigmoidoscopy.

Current methods for alleviating gastrointestinal muscle spasm include the administration of intravenous diazepam, anticholinergics and glucagon, as well as sublingual administration of nitroglycerin. However, these methods produce generalized, systemic effects which persist for a much longer duration than the procedure itself. In addition, nitroglycerin is significantly less effective as a smooth muscle relaxant than S-nitrosothiols, and produces systemic side effects, the most significant of which is hypotension. It is therefore, not used clinically. Clearly, a need exists for a topical smooth muscle relaxant which could be directly instilled into the various regions of the gastrointestinal tract to facilitate both diagnostic and therapeutic endoscopic procedures.

Patient studies, conducted by the inventors, have measured the efficacy of S-nitrosothiols both in facilitating cannulation of the sphincter of Oddi, and in decreasing colon motility to allow for removal of colon polyps. As shown in Figure 1, topical administration of S-nitroso-N-acetylcysteine eliminated duodenal motility. As shown in Figure 2, topical administration of S-nitroso-N-acetylcysteine inhibited the contractile activity of the Sphincter of Oddi, and thus, permitting successful endoscopic cannulation of the sphincter. In addition, administration of S-nitroso-N-acetylcysteine eliminated colon motility to facilitate successful removal of colon polyps. Notably, the relaxant effects were temporary (lasting only for the duration of the procedure), completely reversible and produced no change in systemic blood pressure, heart rate or oxygen saturation. The same type of effects would occur with the use of other cell impermeable S-nitrosothiols, such as S-nitroso-glutathione.

- 17 -

Prior to the present invention, there were no available pharmacological agents which could be applied directly by endoscopic means to exert a direct, immediate, localized, and completely reversible relaxant effect on gastrointestinal smooth muscle. Topical administration of S-nitrosothiols, during endoscopy, eliminates systemic side effects and allows for the use of the lowest effective concentration of the drug.

Administration of S-nitrosothiols obviates the need for sphincterotomy, a procedure which substantially increases the morbidity and mortality of ERCP. In addition, administration of S-nitrosothiols aids in the cannulation and manipulation of the pancreatic duct and biliary tract during therapeutic procedures such as gall bladder cannulation, bile duct stone removal and stent placement, and decreases the incidence of post-ERCP complications, such as pancreatitis and cholangitis. Another use of S-nitrosothiols involves the intraoperative injection of these compounds into the gall bladder prior to cholecystectomy to alleviate cystic duct spasm. This would allow for a laparoscopic cholangiogram by providing access to the cystic duct. In addition to the uses discussed above, S-nitrosothiols may also be administered to treat or prevent any other technical problems associated with endoscopy which are known to those in the medical art.

Another embodiment of the invention relates to administration of a therapeutically effective amount of an S-nitrosothiol compound to relax corpus cavernosum smooth muscle. The term "corpus cavernosum" refers to two areas of smooth muscle which lie side by side on the dorsal aspect of the penis, and together with the corpus spongiosum that surrounds the urethra, constitute erectile tissue. This erectile tissue consists of an irregular sponge-like system of vascular spaces interspersed between arteries and veins. Erection occurs when cavernosa smooth muscle relaxation causes a decrease in arterial resistance and resulting increase in arterial blood flow to the penis.

- 18 -

Smooth muscle has a critical role in erectile function. Thus, another embodiment of the invention relates to the administration of a therapeutically effective amount of an S-nitrosothiol compound for the treatment of impotence. "Impotence" refers to a condition of male sexual dysfunction which is characterized by the inability to obtain or maintain an erection.

Organic causes of erectile impotence, may include endocrine, drug-induced, local injury, neurologic, and vascular. In particular, impotence may result from neurologic blockade caused by such drugs as antihistamines, antihypertensives, psychogenic agents, and anticholinergics. Impotence may also result from neurologic disorders such as anterior temporal lobe lesions, spinal cord disorders, and insufficiency of sensory input resulting from diabetic neuropathy. An additional cause of impotence is insufficient blood flow into the vascular network resulting from an intrinsic defect, or from penile trauma.

Currently available methods for treating impotence consist largely of surgical techniques and intracavernosal injections of pharmacological agents. One surgical technique involves the implantation of a penile prosthesis by inserting within the corpora, a small silastic rod. However, this method does not produce full erection and the complication rate is high. Alternatively, an inflatable prosthetic device may be implanted on either side of the corpora, with a connecting reservoir of material placed in the perivascular space. Erection is achieved through the use of pumps which are located in the scrotum.

Intracavernous injection of the smooth muscle relaxant, papaverine has been used to induce erections. However, a significant disadvantage of this treatment method is the need for a painful injection each time an erection is desired. In addition, numerous side effects and complications result from the chronic use of drugs such as papaverine. Clinical reports indicate that a significant proportion of potential candidates refuse these

- 19 -

injections from the onset of treatment. A larger number of patients, even after favorable initial response to the treatment, become increasingly unresponsive or unwilling to continue injections as a means of treatment (Morales *et al.*, *World J. Urol.* 8:80-83 (1990)).

5 In general, a significant number of patients who are potential candidates for current methods of impotence treatment refuse initially because of the invasive nature of the treatment, or reject further treatment because of pain, fibrosis, or dissatisfaction with results.

10 As demonstrated by the discussion above, prior to the present invention, there was a significant need for a less invasive approach to the treatment of impotence. Because they exert a relaxant effect on corpus cavernosal smooth muscle, S-nitrosothiols are particularly well suited for the treatment of impotence.

15 Administration of S-nitrosothiols results in relaxation of corpus cavernosum smooth muscle, which leads to dilation of the cavernosal arteries and a concomittent increase in blood flow. S-nitrosothiols provide significant advantages in the treatment of impotence over current treatment methods, because they can be administered topically, thereby eliminating the systemic side effects, significant discomfort, fibrosis, and
20 ineffectiveness associated with the currently available, invasive methods of treatment.

Another embodiment of the claimed invention relates to the administration of a therapeutically effective amount of an S-nitrosothiol compound to relax bladder smooth muscle. Bladder smooth muscle
25 includes that of the bladder base, bladder body and proximal urethra. In addition, S-nitrosothiols may be used for the treatment of bladder dysfunction disorders which involve relaxation of bladder smooth muscle. Such disorders include, but are not limited to, problems with bladder filling, volume and continence.

- 20 -

In addition, S-nitrosothiols may be administered to cause relaxation of urethral and bladder base smooth muscle, and thus, facilitate cystoscopic examination of the urinary tract. The term "cystoscopic examination" refers to the introduction of a fiberoptic instrument through the urethra and into the bladder, to achieve visualization of the interior of the urethra and bladder for diagnostic and therapeutic purposes.

Another embodiment of the invention relates to the administration of a therapeutically effective amount of an S-nitrosothiol compound to relax uterine smooth muscle. Increased contractility of uterine smooth muscle precipitates premature labor. Thus, an additional embodiment of the invention relates to the administration of S-nitrosothiol compounds for the treatment or prevention of premature labor.

S-nitrosothiols may also be used to relax fallopian tube smooth muscle. Fallopian tube smooth muscle plays a role in the transport of the egg to the uterus. Thus, S-nitrosothiols may be used to regulate ovum transport, or to facilitate laparoscopic examination of the fallopian tubes, or to facilitate fertilization procedures.

The long-chain lipophilic compounds have unique potential for NO delivery by incorporation into cell membranes, and for accessing the central nervous system (CNS). In the CNS, nitric oxide has been shown to inhibit cell death resulting from ischemic injury, as well as to possess neurotransmitter functions. Membrane permeability achieved by these compounds also provides the unique potential for NO delivery in every organ system. In addition, NO delivery can be regulated by the incorporation of additional functional groups into the molecule. Each functional group, including but not limited to nitrite, nitrate, redox metal, amine, aromatic, and basic amino acids, has its own unique functional aspects which will affect (a) a targeted site of delivery (b) rate of NO release (c) lipophilicity (d) cell permeability (e) duration of action (f) bioactivity and (g) nitrosation potential.

- 21 -

An additional embodiment of the invention relates to the administration of an S-nitrosothiol compound as part of a pharmaceutical composition, comprising a pharmaceutically acceptable carrier, to achieve the physiological effects discussed above.

5 The pharmaceutical compositions utilized in this invention can be administered by intranasal, oral, enteral, topical, sublingual, rectal, intramuscular, intravenous, or subcutaneous means. The compositions may be administered by medical instrumentation including, but not limited to, bronchoscopic, endoscopic, laparoscopic, and cystoscopic means. With
10 respect to the administration of these composition for the treatment of impotence, the term "topical" includes administration in the form of a condom which contains the pharmaceutical composition. Certain S-nitrosothiols, such as lipophilic S-nitrosothiols, are especially suitable for (i.e. lipophilic) incorporation into the condom itself, to provide sustained
15 release of the compound.

 The compounds of this invention can be employed in combination with conventional excipients; i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral or intranasal application which do not deleteriously react with the active compounds.
20 Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethylcellulose, polyvinylpyrrolidone,
25 etc. The pharmaceutical preparations can be sterilized and if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavoring and/or aromatic substances and the like which do not deleteriously react with the active compounds.

For parenteral application, particularly suitable vehicles consist of solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants, including suppositories. Ampules are convenient unit dosages.

5 For enteral application, particularly suitable are tablets, dragees or capsules having talc and/or a carbohydrate carrier binder, or the like, the carrier preferably being lactose and/or corn starch and/or potato starch. A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including
10 those wherein the active component is protected with differentially degradable coatings, e.g., by microencapsulation, multiple coatings, etc.

It will be appreciated that the actually preferred amounts of active compounds used will vary according to the specific compound being utilized, the particular compositions formulated, the mode of application
15 and the particular site of administration. Optimal administration rates for a given protocol of administration can be readily ascertained by those skilled in the art, using conventional dosage determination tests conducted with regard to the foregoing guidelines.

According to the present invention, a "therapeutically effective
20 amount" of a pharmaceutical composition is an amount which is sufficient to achieve the desired pharmacological effect. Generally, the dosage required to provide an effective amount of the composition, and which can be adjusted by one of ordinary skill in the art, will vary, depending upon the age, health, physical condition, sex, weight and extent of disease,
25 of the recipient. Additionally, the dosage may be determined by the frequency of treatment and the nature and scope of the desired effect.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following examples are, therefore, to be construed

- 23 -

as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The entire text of all publications cited above and below are hereby incorporated by reference.

5

EXAMPLES

Example 1. Airway Smooth Muscle Relaxation by S-nitrosothiols

A. Methods

1. Materials

10 Glutathione, L-cysteine, DL-homocysteine, D-penicillin,
 hemoglobin (bovine), methylene blue and Medium 199 sets were
 purchased from Sigma Chemical Co., St. Louis, MO. N-acetylcysteine was
 obtained from Aldrich Chemical Co., Milwaukee, WI. Captopril was
 kindly provided by Dr Victor Dzau. Sodium nitrite, histamine and
 methacholine were purchased from Fisher Scientific, Fairlawn, NJ.
15 Leukotriene D₄ was purchased from Anaquest, BOC Inc., Madison, WI.
 Antibiotic/antimycotic mixture (10,000 U/ml penicillin G sodium, 10,000
 mcg/ml, streptomycin sulfate, 25 mcg/ml amphotericin B) was purchased
 from Gibco Laboratories, Grand Island, NY. Radioimmunoassay kits for
 the determination of cyclic GMP were purchased from New England
20 Nuclear, Boston, MA.

- 24 -

2. Preparation of Airways

Male Hartley guinea pigs (500-600g) were anesthetized by inhalation of enflurane to achieve a surgical plane of anesthesia. The trachea were excised and placed in Kreb's-Henseleit buffer (mM): NaCl 118, CKI 5.4, NaH_2PO_4 1.01, glucose 11.1, NaHCO_3 25.0, MgSO_4 0.69, CaCl 2.32, pH 7.4. The airways were then dissected free from surrounding fat and connective tissue and cut into rings 2-4 mm in diameter. The trachea rings were placed in sterile Medium 199 containing 1% antibiotic/antimycotic mixture in an atmosphere of 5% CO_2 , 45% O_2 , 55% N_2 , and kept for up to 48 hours in tissue culture. The experiments were also performed on human airway smooth muscle, isolated by the same method.

3. Preparation of Blood Vessels

New Zealand White female rabbits weighing 3-4 kg were anesthetized with 30 mg/kg IV sodium pentobarbital. Descending thoracic aortic were isolated and placed immediately in a cold physiologic salt solution (Kreb's) (mM): NaCl 118, CCl 4.7, CaCl_2 2.5, MgSO_4 1.2, KH_2PO_4 1.2, NaHCO_3 12.5, and D-glucose 11.0, pH 7.4. The vessels were cleaned of adherent connective tissue, and the endothelium removed by gentle rubbing with a cotton tipped applicator inserted into the lumen, and cut into 5 mm rings.

4. Preparation of S-nitrosothiols

S-nitrosothiols were prepared at 25°C by reacting equimolar (100 μM) concentrations of reduced thiols with NaNO_2 in 0.5 N HCl (acidified NaNO_2) as described previously (Kowaluk *et al.*, *J. Pharmacol. Exp. Ther.*

- 25 -

256:1256-1264 (1990); Loscalzo *et al.*, *J. Pharmacol. Exp. Ther.* 249(3):726-729 (1989); and Ignarro *et al.*, *J. Pharmacol. Exp. Ther.* 218(3):739-749 (1981)). Solutions turned from clear to various shades of red instantaneously upon product formation, with the notable exception of S-nitroso-penicillamine, which is green.

In this method of synthesis, the reaction of thiols with NO (generated from sodium nitrite) is complete and stoichiometric (Aldred *et al.*, *J. Chem. Soc. Perkin Trans. II*:777-782 (1987); Byler *et al.*, *J. Agric. Food Chem.* 31:523-527 (1983)).

The long-carbon chain lipophilic nitrosothiols, long and short chain S-nitrosodithiols, and S-nitrosothiols with additional functional groups were synthesized by one or more of the following methods: (a) exposure to equimolar N_2O_3 or N_2O_4 in CCl_4 ; (b) exposure to equimolar acidified nitrite; (c) exposure to equimolar bubbled NO gas; (d) exposure to excess cold bubbled NO_2 gas; and (e) exposure to methanolic acid or equimolar $NaNO_2$ diluted in methanol.

The synthesis of S-nitroso-homocysteine has not been previously characterized. This compound displayed the distinct absorption maxima of other S-nitrosothiols at approximately 340 nm and 550 nm (Kowaluk *et al.*, *J. Pharmacol. Exp. Ther.* 256:1256-1264 (1990); Loscalzo *et al.*, *J. Pharmacol. Exp. Ther.* 249(3):726-729 (1989); and Ignarro *et al.*, *J. Pharmacol. Exp. Ther.* 218(3):739-749 (1981)). The molar absorptivity of S-nitroso-homocysteine at 547 nm is $16.7 \text{ cm}^2 \text{M}^{-1}$ and correlates well with published values of 16.6 and 16.1, for S-nitro-cysteine and S-nitro-glutathione, respectively (Kowaluk *et al.*, *J. Pharmacol. Exp. Ther.* 256:1256-1264 (1990)).

Owing to the modest decay of S-nitrosothiols over time, fresh examples were made at hourly intervals and kept at 4°C until use. Solutions were diluted as necessary into physiologic buffer immediately prior to each experiment.

5. Bioassay

Trachea and aortic rings were mounted on stirrups and connected to transducers (model FOT3C Grass) with which changes in isometric tension were measured. Rings were then suspended in 10 cc of oxygenated (95% O₂, 5% CO₂) buffer. Conditions for both the vessel and airway bioassays were established according to standard methodologies as described in Cooke *et al.*, *Am. J. Physiol.* 28:H804-H812 (1989).

In airway experiments, the rings were equilibrated for 60 minutes under a load of 1 gm and then primed twice by exposure to 100 μ M methacholine. Tissues were contracted with various agonists at concentrations determined to generate 50% (\pm 16% S.D.) of maximum tone, after which the effects of different thiols and their S-nitrosylated derivatives were assessed. In selected experiments, relaxation responses were determined in the presence of hemoglobin, or after rings had been preexposed to methylene blue for 30 minutes.

In vessel experiments, aortic rings were contracted with 1 μ M epinephrine and relaxations were induced with S-nitrosothiols.

6. Cyclic Nucleotide Assays

The mechanism by which S-nitrosothiols relax vascular smooth muscle is felt to be through activation of guanylate cyclase with consequent increase in intracellular cyclic GMP (Ignarro *et al.*, *Circ. Res.* 65:1-21 (1989); Loscalzo *et al.*, *J. Pharmacol. Exp. Ther.* 249(3):726-729 (1989)). In order to assess this mechanism in airways, tracheal rings in Krebs's-Henseleit solution were exposed to 100 μ M S-nitroso-N-acetylcysteine (SNOAC) for 90 seconds. Reactions were terminated by

- 27 -

the addition of ice cold 10% trichloroacetic acid and rapid freezing in ethanol-saturated dry ice.

In selected experiments, rings were preexposed to the guanylate cyclase inhibitor, methylene blue (10^{-4} M) for 30 minutes. Tissues were then individually pulverized with a glass (s) homogenizer and centrifuged at 8000 g for 5 minutes. The clear supernatant was extracted with water-saturated ether and assayed for cyclic GMP by radioimmunoassay. Acetylation of samples with acetic anhydride was used to increase the sensitivity of the assay and the determination of recoveries was aided by the use of [3 H] cyclic GMP.

Dose-response relationships to SNOAC were obtained in airways contracted with 3 μ M histamine, and repeated in the presence of 10^{-4} M hemoglobin, 10^{-5} M methylene blue, and 10^{-4} M methylene blue. Relaxation responses to SNOAC are inhibited by hemoglobin and methylene blue, with the latter in a dose-dependent manner. Cyclic GMP determinations were performed in duplicate for each experiment.

7. Statistics

All results are presented as means \pm SEM. Paired samples were compared by the Student's t-test. Dose-response curves were compared by two-way analysis of variance (ANOVA). Values of $p < 0.05$ were considered significant.

- 28 -

Table 1 Inhibitory Concentrations Inducing 50% Relaxation (IC ₅₀)	
<u>RS-NO</u>	<u>IC₅₀ mean ± S.D.; x 10⁻⁶ M</u>
S-nitroso-glutathione	0.99 ± 2.0
S-nitroso-cysteine	3.2 ± 0.2
S-nitroso-homocysteine	2.1 ± 0.3
S-nitroso-N-acetylcysteine	2.1 ± 0.8
S-nitroso-penicillamine	1.8 ± 0.8
S-nitroso-captopril	20.0 ± 0.7

B. Results and Discussion

The mammalian fraction of sulfur that exists as free sulfhydryl is contained largely in the form of glutathione, cysteine, and homocysteine (Jocelyn, P.C., In *Biochemistry of the SH Group*, Academic Press, London/New York pp. 1-46 (1972)). N-acetylcysteine is a minor metabolite of cysteine that is used for its mucolytic properties in the treatment of airway obstruction. N-acetylcysteine has also received attention within the context of nitrate metabolism and undergoes S-nitrosylation in plasma upon treatment with nitroglycerin (Fung *et al.*, *J. Pharmacol. Exp. Ther.* 245(2):524-531 (1988)). The S-nitrosylated derivatives of these four sulfhydryls comprise the group of biological S-nitrosothiols under investigation.

Captopril and penicillamine are examples of nonbiological low molecular weight thiols, and their S-nitrosylated derivatives have been well characterized (Kowaluk *et al.*, *J. Pharmacol. Exp. Ther.* 256:1256-1264

- 29 -

(1990); Loscalzo *et al.*, *J. Pharmacol. Exp. Ther.* 249(3):726-729 (1989); and Ignarro *et al.*, *J. Pharmacol. Exp. Ther.* 218(3):739-749 (1981).

An initial examination of the relaxant activity of each of the biological and nonbiological S-nitrosothiols in guinea pig tracheal rings was conducted. The results are shown in Figures 1 and 2(a)-(f). As demonstrated by dose-response relationships, these compounds are potent airway smooth muscle relaxants, with relaxant effects that are unmatched by equimolar amounts of reactant thiol or NO (generated from NaNO₂ alone).

In every case, the dose-response curves for the S-nitrosothiols were significantly different from the dose-response curves for NO and for the individual thiols by two-way ANOVA to $p < 0.001$. Results are presented as mean \pm SEM, ($n = 5$).

With the exception of S-nitroso-captopril (SNOCAP), the S-nitrosothiols revealed comparable bioactivity with IC₅₀s in the range of 1×10^{-6} M (Table 1). SNOAC and SNOCAP were then selected as representative biological and nonbiological S-nitrosothiols for further detailed investigation.

Dose-effect relationships were obtained for SNOAC and SNOCAP using tracheal rings induced to constrict with leukotriene D₄, histamine, and methacholine. As shown in Figure 3, airways exhibited agonist specificity toward S-nitrosothiol-mediated relaxations: S-nitrosothiols were most active for relaxation of leukotriene D₄-induced contractions and progressively less effective with contractions induced by histamine and methacholine. In every case, SNOAC was approximately 10-fold more active in relaxation of airways than SNOCAP. Results are presented as mean \pm SEM ($n=3-5$).

The time course of relaxation to SNOAC is shown in Figure 4. Using a concentration (5×10^{-6} M) selected to induce approximately 50% of the maximal response, maximal relaxation occurred by five minutes and

- 30 -

a significant residual loss of tone was still evident at one hour. In control experiments, airway contractions remained stable over the study period.

These relaxation responses contrast markedly with those generally ascribed to low-molecular-weight S-nitrosothiols. Figure 5 illustrates the notable difference in relaxation kinetics between these tissues. In vascular smooth muscle, the relaxations are rapid and transient, whereas in tracheal smooth muscle, relaxations occur more slowly and persist for a much longer duration.

Figure 6 shows a comparison between the efficacy of SNOAC and isoproterenol or theophylline under identical study conditions. Of the drugs evaluated, isoproterenol was the most active relaxant, however, SNOAC was approximately 50 times more active in relaxation than theophylline. The dose response curves for these agents are each significantly different from each other by two-way ANOVA to $p < 0.01$. Results are expressed as mean \pm SEM ($n=3-5$).

Hemoglobin and methylene blue are established inhibitors of NO-induced relaxations in vascular smooth muscle (Ignarro *et al.*, *Circ. Res.* 65:1-21 (1989)). When their effects were examined in airways, hemoglobin and methylene blue each significantly attenuated the actions of SNOAC, as evidenced by rightward shifts in the dose-effect relationships to SNOAC in their presence (Figure 7). The dose-response curves for SNOAC were significantly different from each of the curves derived in the presence of hemoglobin and methylene blue by two-way ANOVA to $p=0.05$. Results are presented as mean \pm SEM ($n=3-5$).

The biochemical mechanism of action of S-nitrosothiols was further investigated in isolated tracheal rings. As shown in Figure 8, tracheal rings incubated with SNOAC exhibited 4-fold increases in cyclic GMP over basal levels (control). Increases in cyclic GMP were attenuated by pretreatment of tissues with the guanylate cyclase inhibitor, methylene blue (10^{-4} M). Cyclic GMP levels in the presence of SNOAC were

- 31 -

significantly greater than control values ($p < 0.0005$) and levels determined in the presence of methylene blue ($p = 0.05$). Results are presented as mean \pm SEM ($n = 4-8$).

An examination of the relaxant activity of S-nitrosothiols in human tracheal rings was also conducted. The results are shown in Figures 11-15. In particular, Figure 11 shows that S-nitroso-glutathione has a relaxant effect upon human trachea which is significantly greater than nitrite (NO_2). Figure 12 demonstrates that the relaxant effect of S-nitrosoglutathione upon human trachea is significantly greater than glutathione alone. This data underscores the fact that the bioactivity of nitric oxide in airways depends upon the form in which it is delivered. S-nitrosothiols provide efficient delivery of NO in its most bioactive form.

Figure 13 demonstrates that the relaxant effect of SNOAC upon human trachea is significantly greater than that of N-acetylcysteine. As shown in Figure 13, NAC caused significant constriction of the tracheal smooth muscle, which is consistent with the fact that NAC, when given as a mucolytic agent, causes the untoward side effect of bronchospasm. SNOAC not only causes relaxation of airway tissue, but also eliminates bronchospasm.

Figure 14 demonstrates that SNOAC and SNOGSH exert a relaxant effect on airway smooth muscle which is significantly more potent than that of theophylline, and compares favorably with that exerted by isoproterenol.

Experiments were also conducted to assess the cGMP response to SNOAC in human airways. As shown in Figure 15, tracheal rings incubated with SNOAC exhibited 4-fold increases in cyclic GMP over basal levels (control).

Unexpectedly, the relaxation response to low molecular weight S-nitrosothiols in airways differs markedly from that observed in blood vessels. In the latter case, relaxations occur slowly and persist for a much

longer duration. This is most likely attributed to the inherent differences between vascular and nonvascular smooth muscle. There may be additional contributing factors responsible for this heterogeneity. Finally, any disparity among smooth muscle cells in redox state, availability of
5 reducing equivalents, pH, oxygen tension, or any other factor that might influence the stability of the S-NO bond would be predicted to influence relaxation kinetics.

The results also suggests that, in addition to the primary site of obstruction in the lung, the efficacy of nitro(so)-bronchodilators may be
10 determined by the nature of the chemical mediators contributing to bronchoconstriction. In particular, S-nitrosothiols were most effective in this study against leukotriene D₄-mediated bronchoconstriction and progressively less effective against histamine and methacholine-mediated constriction. Thus, regional variation in guanylate cyclase content or
15 activity, the site of obstruction, the form in which the active species of NO is administered, and the nature of the bronchoconstrictor stimuli are all variables which may influence the determination of nitro(so)-bronchodilator efficacy and the importance of guanylate cyclase in regulating airway tone.

20 In summary, S-nitrosothiols are believed to be important intermediates in the metabolism of organic nitrates and endogenously-derived NO. These compounds are more stable than NO and retain its cyclic GMP-dependent bioactivity in blood vessels.

In the present invention, the inventors have demonstrated that S-
25 nitrosothiols are also potent airway smooth muscle relaxants and mediate their effects through activation of guanylate cyclase. The results indicate that cyclic GMP is an important inhibitory mediator of airway tone, and suggest a mechanism by which the bioactivity of NO is preserved in the presence of high, ambient concentrations of oxygen and oxygen-derived
30 free radicals. The inventors have also demonstrated that S-nitrosothiols


- 33 -

are present in human airway secretions and mediate the airway relaxation caused by VIP. As a result of their potent relaxation effect upon airway smooth muscle S-nitrosothiols have significant pharmacological utility for the treatment of airway obstruction.

- 34 -

WHAT IS CLAIMED IS:


1. A compound having the formula:



wherein:

X equals 2 to 20.


2. A compound having the formula:



wherein:

X equals 2 to 20.

3. A compound having the formula:



wherein:

X equals 2 to 20;


Y is selected from the group consisting of fluoro, C₁-C₆ alkoxy, cyano, carboxamido, C₃-C₆ cycloalkyl, aralkoxy, C₂-C₆ alkylsulfinyl, arylthio, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbomoyl, C₁-C₆ N-alkylcarbomoyl, C₂-C₁₅ N,N-dialkylcarbomoyl, amino, hydroxyl, carboxyl, hydrogen, nitro and aryl;

wherein aryl includes benzyl, naphthyl and anthracenyl groups.

4. A method for relaxing airway smooth muscle, comprising administering a therapeutically effective amount of an S-nitrosothiol compound to an animal in need thereof.

5. The method of claim 4 wherein said S-nitrosothiol compound has the formula:


- 35 -



wherein:

X equals 2 to 20.


6. The method of claim 4 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.

7. The method of claim 4 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20;

Y is selected from the group consisting of fluoro, $\text{C}_1\text{-C}_6$ alkoxy, cyano, carboxamido, $\text{C}_3\text{-C}_6$ cycloalkyl, aralkoxy, $\text{C}_2\text{-C}_6$ alkylsulfinyl, arylthio, $\text{C}_1\text{-C}_6$ alkylamino, $\text{C}_2\text{-C}_{15}$ dialkylamino, hydroxy, carbomoyl, C_1C_6 N-alkylcarbomoyl, $\text{C}_2\text{-C}_{15}$ N,N-dialkylcarbomoyl, amino, hydroxyl, carboxyl, hydrogen, nitro and aryl;

wherein aryl includes benzyl, naphthyl and anthracenyl groups.

8. The method of claim 4 wherein said S-nitrosothiol compound is selected from the group consisting of S-nitroso-N-acetylcysteine, S-nitroso-glutathione, S-nitroso-cysteine, S-nitroso-homocysteine, S-nitroso-penicillamine and S-nitroso-captopril.


- 36 -

9. The method of claim 4 wherein said compound is administered as part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

10. The method of claim 9 wherein said pharmaceutical composition is administered to an animal by a route comprising oral, sublingual, intravenous, intramuscular, aerosol, topical or bronchoscopic delivery.

11. A method for treatment or prevention of respiratory disorders, comprising administering a therapeutically effective amount of an S-nitrosothiol compound to an animal in need thereof.


12. The method of claim 11 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.


13. The method of claim 11 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.

14. The method of claim 11 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20;

- 37 -

Y is selected from the group consisting of fluoro, C₁-C₆ alkoxy, cyano, carboxamido, C₃-C₆ cycloalkyl, aralkoxy, C₂-C₆ alkylsulfinyl, arylthio, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbomoyl, C₁C₆ N-alkylcarbomoyl, C₂-C₁₅ N,N-dialkylcarbomoyl, amino, hydroxyl, carboxyl, hydrogen, nitro and aryl;

wherein aryl includes benzyl, naphthyl and anthracenyl groups.

15. The method of claim 11 wherein said S-nitrosothiol compound is selected from the group consisting of S-nitroso-N-acetylcysteine, S-nitroso-glutathione, S-nitroso-cysteine, S-nitroso-homocysteine, S-nitroso-penicillamine and S-nitroso-captopril.

16. The method of claim 11 wherein said compound is administered as part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.


17. The method of claim 16 wherein said pharmaceutical composition is administered to an animal by a route comprising oral, sublingual, intravenous, intramuscular, aerosol topical or bronchoscopic delivery.

18. The method of claim 11 wherein said respiratory disorder is in the group comprised of all subsets of obstructive lung disease, including emphysema, asthma, bronchitis, fibrosis, excessive mucous secretion, obstruction of air flow, and lung disorders resulting from post-surgical complications.

- 38 -

19. A method for relaxing gastrointestinal smooth muscle, comprising administering a therapeutically effective amount of an S-nitrosothiol compound to an animal in need thereof.


20. The method of claim 19 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.


21. The method of claim 19 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.

22. The method of claim 19 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20;

Y is selected from the group consisting of fluoro, C₁-C₆ alkoxy, cyano, carboxamido, C₃-C₆ cycloalkyl, aralkoxy, C₂-C₆ alkylsulfinyl, arylthio, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbomoyl, C₁C₆ N-alkylcarbomoyl, C₂-C₁₅ N,N-dialkylcarbomoyl, amino, hydroxyl, carboxyl, hydrogen, nitro and aryl;

wherein aryl includes benzyl, naphthyl and anthracenyl groups.

- 39 -


23. The method of claim 19 wherein said S-nitrosothiol compound is selected from the group consisting of S-nitroso-N-acetylcysteine, S-nitroso-glutathione, S-nitroso-cysteine, S-nitroso-homocysteine, S-nitroso-penicillamine and S-nitroso-captopril.

24. The method of claim 19 wherein said compound is administered as part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

25. The method of claim 24 wherein said pharmaceutical composition is administered to a patient by a route comprising oral, sublingual, intravenous, topical, intramuscular, aerosol or endoscopic delivery.

26. A method for alleviating contraction or spasm of gastrointestinal smooth muscle associated with endoscopic procedures comprising, administering a therapeutically effective amount of an S-nitrosothiol compound to an animal in need thereof.


27. The method of claim 26 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.

28. The method of claim 26 wherein said S-nitrosothiol compound has the formula:




wherein:

X equals 2 to 20.

- 40 -

29. The method of claim 26 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20;

Y is selected from the group consisting of fluoro, C₁-C₆ alkoxy, cyano, carboxamido, C₃-C₆ cycloalkyl, aralkoxy, C₂-C₆ alkylsulfinyl, arylthio, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbomoyl, C₁C₆ N-alkylcarbomoyl, C₂-C₁₅ N,N-dialkylcarbomoyl, amino, hydroxyl, carboxyl, hydrogen, nitro and aryl;

wherein aryl includes benzyl, naphthyl and anthracenyl groups.

30. The method of claim 26 wherein said S-nitrosothiol compound is selected from the group consisting of S-nitroso-N-acetylcysteine, S-nitroso-glutathione, S-nitroso-cysteine, S-nitroso-homocysteine, S-nitroso-penicillamine and S-nitroso-captopril.


31. The method of claim 26 wherein said compound is administered as part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

32. The method of claim 31 wherein said pharmaceutical composition is administered to an animal by a route comprising oral, sublingual, intravenous, topical, intramuscular, aerosol or endoscopic delivery.

33. A method for relaxing corpus cavernosum smooth muscle, comprising administering a therapeutically effective amount of an S-nitrosothiol compound to an animal in need thereof.

- 41 -


34. The method of claim 33 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.


35. The method of claim 33 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.

36. The method of claim 33 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20;

Y is selected from the group consisting of fluoro, C₁-C₆ alkoxy, cyano, carboxamido, C₃-C₆ cycloalkyl, aralkoxy, C₂-C₆ alkylsulfinyl, arylthio, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbomoyl, C₁C₆ N-alkylcarbomoyl, C₂-C₁₅ N,N-dialkylcarbomoyl, amino, hydroxyl, carboxyl, hydrogen, nitro and aryl;

wherein aryl includes benzyl, naphthyl and anthracenyl groups.

37. The method of claim 33 wherein said S-nitrosothiol compound is selected from the group consisting of S-nitroso-N-acetylcysteine, S-nitroso-glutathione, S-nitroso-cysteine, S-nitroso-homocysteine, S-nitroso-penicillamine and S-nitroso-captopril.


- 42 -

38. The method of claim 33 wherein said compound is administered as part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

39. The method of claim 38 wherein said pharmaceutical composition is administered to an animal by a route comprising oral, sublingual, intravenous, topical, intramuscular or aerosol delivery.

40. A method for the treatment or prevention of impotence, comprising administering a therapeutically effective amount of an S-nitrosothiol compound to an animal in need thereof.


41. The method of claim 40 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.


42. The method of claim 40 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.

43. The method of claim 40 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20;

- 43 -

Y is selected from the group consisting of fluoro, C₁-C₆ alkoxy, cyano, carboxamido, C₃-C₆ cycloalkyl, aralkoxy, C₂-C₆ alkylsulfinyl, arylthio, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbomoyl, C₁C₆ N-alkylcarbomoyl, C₂-C₁₅ N,N-dialkylcarbomoyl, amino, hydroxyl, carboxyl, hydrogen, nitro and aryl;

wherein aryl includes benzyl, naphthyl and anthracenyl groups.


44. The method of claim 40 wherein said S-nitrosothiol compound is selected from the group consisting of S-nitroso-N-acetylcysteine, S-nitroso-glutathione, S-nitroso-cysteine, S-nitroso-homocysteine, S-nitroso-penicillamine and S-nitroso-captopril.

45. The method of claim 40 wherein said compound is administered as part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

46. The method of claim 45 wherein said pharmaceutical composition is administered to an animal by a route comprising oral, sublingual, intravenous, topical, intramuscular or aerosol delivery.

47. A method for relaxing bladder smooth muscle, comprising administering a therapeutically effective amount of an S-nitrosothiol compound to an animal in need thereof.

48. The method of claim 47 wherein said S-nitrosothiol compound has the formula:




wherein:

X equals 2 to 20.

- 44 -


49. The method of claim 47 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.

50. The method of claim 47 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20;

Y is selected from the group consisting of fluoro, $\text{C}_1\text{-C}_6$ alkoxy, cyano, carboxamido, $\text{C}_3\text{-C}_6$ cycloalkyl, aralkoxy, $\text{C}_2\text{-C}_6$ alkylsulfinyl, arylthio, $\text{C}_1\text{-C}_6$ alkylamino, $\text{C}_2\text{-C}_{15}$ dialkylamino, hydroxy, carbomoyl, C_1C_6 N-alkylcarbomoyl, $\text{C}_2\text{-C}_{15}$ N,N-dialkylcarbomoyl, amino, hydroxyl, carboxyl, hydrogen, nitro and aryl;

wherein aryl includes benzyl, naphthyl and anthracenyl groups.

51. The method of claim 47 wherein said S-nitrosothiol compound is selected from the group consisting of S-nitroso-N-acetylcysteine, S-nitroso-glutathione, S-nitroso-cysteine, S-nitroso-homocysteine, S-nitroso-penicillamine and S-nitroso-captopril.


52. The method of claim 47 wherein said compound is administered as part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

- 45 -

53. The method of claim 52 wherein said pharmaceutical composition is administered to an animal by a route comprising oral, sublingual, intravenous, topical, intramuscular or aerosol delivery.

54. A method for relaxing uterine smooth muscle, comprising administering a therapeutically effective amount of an S-nitrosothiol compound to an animal in need thereof.


55. The method of claim 54 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.


56. The method of claim 54 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20.

57. The method of claim 54 wherein said S-nitrosothiol compound has the formula:



wherein:

X equals 2 to 20;

Y is selected from the group consisting of fluoro, C₁-C₆ alkoxy, cyano, carboxamido, C₃-C₆ cycloalkyl, aralkoxy, C₂-C₆ alkylsulfinyl, arylthio, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbomoyl, C₁C₆ N-alkylcarbomoyl, C₂-C₁₅ N,N-dialkylcarbomoyl, amino, hydroxyl, carboxyl, hydrogen, nitro and aryl;

- 46 -

wherein aryl includes benzyl, naphthyl and anthracenyl groups.

58. The method of claim 54 wherein said S-nitrosothiol compound is selected from the group consisting of S-nitroso-N-acetylcysteine, S-nitroso-glutathione, S-nitroso-cysteine, S-nitroso-homocysteine, S-nitroso-penicillamine and S-nitroso-captopril.

59. The method of claim 54 wherein said compound is administered as part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

60. The method of claim 59 wherein said pharmaceutical composition is administered to an animal by a route comprising oral, sublingual, intravenous, topical, intramuscular or aerosol delivery.